

Big ideas from last time

Rather than measuring the “size” of \mathcal{H} with $|\mathcal{H}|$, we can instead think about:

Using \mathcal{H} , how many ways can we label a dataset?

We call a particular labeling of $\mathbf{x}_1, \dots, \mathbf{x}_n$ a **dichotomy**

Using this language, we can answer our question via the growth function $m_{\mathcal{H}}(n)$, which counts the **most** dichotomies that \mathcal{H} could ever generate on n points

It is easy to see that $m_{\mathcal{H}}(n) \leq 2^n$

- If $m_{\mathcal{H}}(n) = 2^n$, we say that \mathcal{H} can **shatter** a set of size n
- If no set of size k can be shattered by \mathcal{H} ($m_{\mathcal{H}}(k) < 2^k$) then k is a **break point**

Goal for today

Using Hoeffding's inequality together with a union bound, we were able to show that

$$\mathbb{P} \left[\max_{h \in \mathcal{H}} |\hat{R}_n(h) - R(h)| > \epsilon \right] \leq |\mathcal{H}| \cdot 2e^{-2\epsilon^2 n}$$

What the VC bound gives us is a generalization of the form

$$\mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - R(h)| > \epsilon \right] \leq 2 \cdot m_{\mathcal{H}}(2n) \cdot 2e^{-\frac{1}{8}\epsilon^2 n}$$

supremum: maximum over an infinite set

Supremum

The **supremum** of a set $S \subset T$ is the least element of T that is greater than or equal to all elements of S

Sometimes called the **least upper bound**

Examples

- $\sup\{1, 2, 3\} = 3$
- $\sup\{x : 0 \leq x \leq 1\} = 1$
- $\sup\{x : 0 < x < 1\} = 1$
- $\sup\{1 - 1/n : n > 0\} = 1$

Why might this work?

We aim to get a bound on $\mathbb{P}[|\hat{R}_n(h) - R(h)| > \epsilon]$ that holds for any $h \in \mathcal{H}$, i.e., a bound on

$$\mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - R(h)| > \epsilon \right] \leftarrow$$

It should not be a shock that we can bound this using the growth function...

There may be infinitely many $h \in \mathcal{H}$, but \mathcal{H} can only generate $m_{\mathcal{H}}(n)$ **unique dichotomies**


Thus, $\hat{R}_n(h)$ can only take at most $m_{\mathcal{H}}(n)$ different values

Unfortunately, $R(h)$ can still take infinitely many different values, and so there are infinitely many $|\hat{R}_n(h) - R(h)|$

Fundamental insight

The key insight (or trick) is to consider *two* datasets!

We will imagine that in addition to our training data, we have access to a second independent dataset (of size n), which we call the *ghost dataset*



Can we relate $\mathbb{P}[|\hat{R}_n(h) - R(h)| > \epsilon]$ to $\mathbb{P}[|\hat{R}_n(h) - \hat{R}'_n(h)| > \epsilon]$?

Using the ghost dataset

Suppose (for the moment) that the empirical estimates $\hat{R}_n(h)$ and $\hat{R}'_n(h)$ are random variables that are drawn from a *symmetric* distribution with mean (and median) $R(h)$

Consider the following events:

- A : the event that $|\hat{R}_n(h) - R(h)| > \epsilon$
- B : the event that $|\hat{R}_n(h) - \hat{R}'_n(h)| > \epsilon$

Claim: $\mathbb{P}[B|A] \geq \frac{1}{2}$

Thus $\mathbb{P}[B] \geq \mathbb{P}[B|A] \cdot \mathbb{P}[A] \geq \frac{1}{2}\mathbb{P}[A]$

$$\longrightarrow \underbrace{\mathbb{P}[|\hat{R}_n(h) - R(h)| > \epsilon]}_A \leq 2 \underbrace{\mathbb{P}[|\hat{R}_n(h) - \hat{R}'_n(h)| > \epsilon]}_B$$

Using the ghost dataset

Unfortunately, the distribution of $\hat{R}_n(h)$ and $\hat{R}'_n(h)$ is binomial (not symmetric) so this exact statement doesn't hold in general, but the intuition is valid

Instead, we have the following bound:

Lemma 1 (Ghost dataset)

$$\mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - R(h)| > \epsilon \right]$$



$$\leq 2 \mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \right]$$



Bounding the worst-case deviation

Lemma 2 (Where the magic happens)

Let $S = \{(x_i, y_i), i = 1, \dots, 2n\}$. Then

$$\begin{aligned} & \mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \right] \\ & \leq m_{\mathcal{H}}(2n) \cdot \underbrace{\sup_S}_{\text{dataset } S \text{ fixed}} \underbrace{\sup_{h \in \mathcal{H}} \mathbb{P} \left[|\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \mid S \right]}_{\text{probability w.r.t. random partition of } S} \end{aligned}$$

here, the dataset S is fixed
the probability is with respect
to a random partition of S into
two training sets of size n

Proof of Lemma 2

It is straightforward to show that

$$\begin{aligned} \rightarrow) \quad & \mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \right] \leftarrow \\ & \leq \sup_S \mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \middle| S \right] \leftarrow \end{aligned}$$

Note that in the probability on the right-hand side, the dataset S is fixed

Thus, there are only a finite number of dichotomies that \mathcal{H} can generate on S

Call this number $m_{\mathcal{H}}(S)$

Let $\underline{h_1, \dots, h_{m_{\mathcal{H}}(S)}}$ be classifiers giving rise to these dichotomies

Proof of Lemma 2

Using this observation, we have

$$\begin{aligned} & \mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \middle| S \right] \\ &= \mathbb{P} \left[\max_{h_1, \dots, h_{m_{\mathcal{H}}(S)}} |\hat{R}_n(h_i) - \hat{R}'_n(h_i)| > \frac{\epsilon}{2} \middle| S \right] \\ &\leq \sum_{i=1}^{m_{\mathcal{H}}(S)} \mathbb{P} \left[|\hat{R}_n(h_i) - \hat{R}'_n(h_i)| > \frac{\epsilon}{2} \middle| S \right] \\ &\leq m_{\mathcal{H}}(S) \max_{h_1, \dots, h_{m_{\mathcal{H}}(S)}} \mathbb{P} \left[|\hat{R}_n(h_i) - \hat{R}'_n(h_i)| > \frac{\epsilon}{2} \middle| S \right] \\ &\leq m_{\mathcal{H}}(2n) \cdot \sup_{h \in \mathcal{H}} \mathbb{P} \left[|\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \middle| S \right] \end{aligned}$$

Final step

At this point, we have shown

$$\mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - R(h)| > \epsilon \right] \\ \rightarrow \leq 2 \cdot m_{\mathcal{H}}(2n) \cdot \sup_S \sup_{h \in \mathcal{H}} \mathbb{P} \left[|\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \middle| S \right]$$

Lemma 3 (Random partitions)

For *any* h and *any* S ,

$$\mathbb{P} \left[|\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \middle| S \right] \leq 2e^{-\frac{1}{8}\epsilon^2 n}$$

Proof follows from a simple lemma (also by Hoeffding)

Putting it all together

$$\begin{aligned} & \mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - R(h)| > \epsilon \right] \\ & \leq 2 \mathbb{P} \left[\sup_{h \in \mathcal{H}} |\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \right] \\ & \leq 2 \cdot m_{\mathcal{H}}(2n) \cdot \sup_S \sup_{h \in \mathcal{H}} \mathbb{P} \left[|\hat{R}_n(h) - \hat{R}'_n(h)| > \frac{\epsilon}{2} \middle| S \right] \\ & \leq 2 \cdot m_{\mathcal{H}}(2n) \cdot 2e^{-\frac{1}{8}\epsilon^2 n} = \delta \end{aligned}$$

Thus, for any $h \in \mathcal{H}$, we have that with probability $\geq 1 - \delta$

$$R(h) \leq \hat{R}_n(h) + \sqrt{\frac{8}{n} \log \frac{4m_{\mathcal{H}}(2n)}{\delta}}$$

Using the VC bound: The VC dimension

We went to a lot of trouble to show that if k is a break point for \mathcal{H} , then

$$m_{\mathcal{H}}(n) \leq \sum_{i=0}^{k-1} \binom{n}{i} \leq n^{k-1} + 1$$

$$\begin{aligned} \Rightarrow R(h) &\leq \hat{R}_n(h) + \sqrt{\frac{8}{n} \log \frac{4((2n)^{k-1} + 1)}{\delta}} \\ &\lesssim \hat{R}_n(h) + \sqrt{\frac{8(k-1)}{n} \log \frac{8n}{\delta}} \end{aligned}$$

True for $k \geq 3$

The **VC dimension** of a hypothesis set \mathcal{H} , denoted $d_{\text{VC}}(\mathcal{H})$, is the largest n for which $m_{\mathcal{H}}(n) = 2^n$

- $d_{\text{VC}}(\mathcal{H})$ is the most points that \mathcal{H} can shatter
- $d_{\text{VC}}(\mathcal{H})$ is 1 less than the smallest break point



$$\Rightarrow R(h) \lesssim \hat{R}_n(h) + \sqrt{\frac{8d_{\text{VC}}}{n} \log \frac{8n}{\delta}}$$

Examples

- Positive rays: $d_{VC} = 1$
- Positive intervals: $d_{VC} = 2$
- Convex sets: $d_{VC} = \infty$
- Linear classifiers in \mathbb{R}^2 : $d_{VC} = 3$

VC dimension of general linear classifiers

For $d = 2$, $d_{VC} = 3$



In general $d_{VC} = d + 1$

We will prove this by showing that $d_{VC} \leq d + 1$ and $d_{VC} \geq d + 1$

One direction

Let's first show that there exists a set of $d + 1$ points in \mathbb{R}^d that are shattered

Consider the matrix

$$\mathbf{X} = \underbrace{\begin{bmatrix} -\tilde{\mathbf{x}}_1^T - \\ -\tilde{\mathbf{x}}_2^T - \\ \vdots \\ -\tilde{\mathbf{x}}_{d+1}^T - \end{bmatrix}}_{d+1} = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & 1 & 0 & \cdots & 0 \\ 1 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \cdots & 1 \end{bmatrix} \begin{bmatrix} b \\ w \end{bmatrix}$$

$X\theta = y$
 $\theta = X^{-1}y$

Think of each row as the concatenation $\tilde{\mathbf{x}}_i^T = [\underline{1}, \underline{\mathbf{x}}_i^T]$

One can show that \mathbf{X} is invertible

$$\begin{aligned} & \langle w, x_i \rangle + b \\ &= \theta^T \tilde{x}_i \\ & \theta = \underline{[b, w^T]} \end{aligned}$$

Can we shatter this data set?

For any $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{d+1} \end{bmatrix} = \begin{bmatrix} \pm 1 \\ \pm 1 \\ \vdots \\ \pm 1 \end{bmatrix}$, can we find a vector $\boldsymbol{\theta}$

satisfying $\text{sign}(\underline{\mathbf{X}\boldsymbol{\theta}}) = \mathbf{y}$?

Easy! Just make $\boldsymbol{\theta} = \mathbf{X}^{-1}\mathbf{y}$ and we have


$$\mathbf{X}\boldsymbol{\theta} = \mathbf{y}$$

$$\text{sign}(\mathbf{X}\boldsymbol{\theta}) = \text{sign}(\mathbf{y}) = \mathbf{y}$$

We can shatter a set of $d + 1$ points

What does this prove?

a) $d_{VC} = d + 1$


b) $d_{VC} \geq d + 1$ 

c) $d_{VC} \leq d + 1$

d) None of the above

To finish the proof

In order to show that $d_{VC} \leq d + 1$, we need to show

- a) There are $d + 1$ points we cannot shatter
- b) There are $d + 2$ points we cannot shatter
- c) We cannot shatter any set of $d + 1$ points
- d) We cannot shatter any set of $d + 2$ points 

The other direction

Take any $d + 2$ points $\tilde{\mathbf{x}}_1, \dots, \tilde{\mathbf{x}}_{d+2}$

We have more points than dimensions, so there must be some j for which

$$\tilde{\mathbf{x}}_j = \sum_{i \neq j} \alpha_i \tilde{\mathbf{x}}_i$$

where not all $\alpha_i = 0$

Consider the dichotomy where the $\tilde{\mathbf{x}}_i$ with $\alpha_i \neq 0$ are labeled $y_i = \text{sign}(\alpha_i)$, and $y_j = -1$

No linear classifier can implement such a dichotomy!

Why not?

$$\tilde{\mathbf{x}}_j = \sum_{i \neq j} \alpha_i \tilde{\mathbf{x}}_i \quad \rightarrow \quad \boldsymbol{\theta}^T \tilde{\mathbf{x}}_j = \sum_{i \neq j} \alpha_i \boldsymbol{\theta}^T \tilde{\mathbf{x}}_i$$

If $y_i = \text{sign}(\boldsymbol{\theta}^T \tilde{\mathbf{x}}_i) = \text{sign}(\alpha_i)$, then $\alpha_i \boldsymbol{\theta}^T \tilde{\mathbf{x}}_i > 0$

This means that $\boldsymbol{\theta}^T \tilde{\mathbf{x}}_j = \sum_{i \neq j} \alpha_i \boldsymbol{\theta}^T \tilde{\mathbf{x}}_i > 0$

Thus $y_j = \text{sign}(\boldsymbol{\theta}^T \tilde{\mathbf{x}}_j) = +1$

Interpreting the VC dimension

We have just shown that for a linear classifier in \mathbb{R}^d

$$\begin{array}{l} d_{VC} \geq d + 1 \\ d_{VC} \leq d + 1 \end{array} \quad \longrightarrow \quad d_{VC} = d + 1$$

How many parameters does a linear classifier in \mathbb{R}^d have?

$$\begin{array}{l} \mathbf{w} \in \mathbb{R}^d \\ b \in \mathbb{R} \end{array} \quad \longrightarrow \quad d + 1$$

The usual examples

- Positive rays
 - $d_{VC} = 1$
 - 1 parameter
- Positive intervals
 - $d_{VC} = 2$
 - 2 parameters
- Convex sets
 - $d_{VC} = \infty$
 - as many parameters as you want

Effective number of parameters

Additional parameters do not always contribute additional degrees of freedom


Example

Take the output of a linear classifier, and then feed this into another linear classifier

$$y_i = \text{sign} (w' (\text{sign}(\mathbf{w}^T \mathbf{x}_i + b)) + b')$$

The parameters w' and b' are totally redundant
(they do not allow us to create any new classifiers/dichotomies)

Interpreting the VC bound



VC bound in action


How big does our training set need to be?

$$R(h) \lesssim \hat{R}_n(h) + \underbrace{\sqrt{\frac{8d_{VC}}{n} \log \frac{8n}{\delta}}}_{\epsilon}$$

Just to see how this behaves, let's ignore the constants and suppose that

$$\epsilon \sim \sqrt{\frac{d_{VC}}{n} \log n}$$

VC bound in action



RULE OF THUMB: $n \geq 10d_{VC}$